

REVIEW

Open Access



Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in West and Central Africa

Emilie Hendrickx¹, Lian F. Thomas^{2,3}, Pierre Dorny^{1,4}, Branko Bobić⁵, Uffe Christian Braae^{6,7}, Brecht Devleeschauwer^{8,9}, Ramon M. Eichenberger¹⁰, Sarah Gabriël⁸, Anastasios Saratsis¹¹, Paul R. Torgerson¹⁰, Lucy J. Robertson¹² and Veronique Dermauw^{1*} 

Abstract

Background: The zoonotic tapeworm *Taenia saginata*, although causing only minor discomfort in humans, is responsible for considerable economic losses in the livestock sector due to condemnation or downgrading of infected beef carcasses. An overview of current knowledge on the distribution and prevalence of this parasite in West and Central Africa is lacking.

Methods: We conducted a systematic review, collecting information on published and grey literature about *T. saginata* taeniosis and bovine cysticercosis from 27 countries/territories in West and Central Africa, published between January 1st, 1990 and December 31st, 2017.

Results: The literature search retrieved 1672 records, of which 51 and 45 were retained for a qualitative and quantitative synthesis, respectively. Non-specified human taeniosis cases were described for Nigeria, Cameroon, Senegal, Burkina Faso, Democratic Republic Congo, Guinea, and Ivory Coast (seven out of 27 countries/territories), while *T. saginata* taeniosis specifically was only reported for Cameroon. Most prevalence estimates for taeniosis ranged between 0–11%, while three studies from Nigeria reported prevalence estimates ranging between 23–50%. None of the studies included molecular confirmation of the causative species. The presence of bovine cysticercosis was reported for Benin, Burkina Faso, Cameroon, Central African Republic, Chad, Democratic Republic Congo, Ghana, Guinea, Ivory Coast, Mali, Niger, Nigeria, Senegal, and Tristan da Cunha (14 out of 27 countries/territories). Prevalence estimates ranged between 0–29%.

Conclusions: Our systematic review has revealed that human taeniosis and bovine cysticercosis are seriously understudied in West and Central Africa. The high prevalence estimates of both conditions suggest an active dissemination of this parasite in the region, calling for a concerted One Health action from public health, veterinary health and food surveillance sectors.

Keywords: *Taenia saginata*, Cysticercosis, Cattle, Beef, Tapeworm, West Africa, Central Africa

*Correspondence: vdermauw@itg.be

¹ Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

Full list of author information is available at the end of the article



Background

The tapeworm *Taenia saginata* is one of three *Taenia* species that infect humans as their definitive host, with bovines serving as the intermediate host. Humans acquire *T. saginata* infection after consuming undercooked beef containing viable cysticerci. The adult tapeworm resides in the small intestine, where it becomes patent within approximately ten weeks. At that moment, the strobila may have reached a length of up to three meters [1], and gravid proglottids can contain up to 100,000 taeniid eggs. These eggs are voided during and between defecation [2], and have the potential to survive for a long time without hatching. Eggs found in faecal material and eggs within soil have been documented to remain viable for up to 9.5 months [3]. Contaminated pastures, water and feed are a source of infection for cattle. Following ingestion, the early larval stages (oncospheres) hatch and the hexacanth larvae migrate, utilising the lymphatic and blood system, to the muscle tissue. Here the larvae mature into the metacestode stage, called cysticerci [4].

Unlike *Taenia solium*, for which humans can also act as a dead-end intermediate host leading to the debilitating and stigmatising disease neurocysticercosis, human *T. saginata* infections are restricted to the definitive (adult tapeworm) stage, which has a more limited public health burden. *Taenia saginata* taeniosis is generally asymptomatic or associated with mild abdominal discomfort, although more serious complications, including appendicitis, intestinal obstruction and gall bladder perforation have occasionally been documented [3]. Bovine cysticercosis, however, may result in substantial economic losses generated for the food industry because of meat condemnation, treatment processing costs and an overall reduction in the product value [5]. Moreover, the meat inspection process itself requires substantial (veterinary) public health sector investment and there are costs associated with treatment seeking behaviour, diagnostics, and treatment of human taeniosis cases [5, 6].

Taenia saginata is considered to have a global distribution, with higher prevalences in low-income regions where sanitation standards may be poor, and the meat inspectorate services are often poorly funded and understaffed.

In West and Central Africa, the cattle population amounts to 120 million heads [7]. While West Africa mainly consists of arid (and to lesser degree semi-arid and sub-humid) agro-ecological zones, Central Africa predominantly consists of humid zones (with some sub-humid zones as well) [8]. In the purely humid agro-ecological zones, cattle production is not considered an important economic activity due to the presence of diseases such as trypanosomiasis [8]. In the arid zones, pastoralism is the most commonly cattle production system,

while the semi-arid and sub-humid zones in the area are characterized by mixed crop-livestock farming systems [8].

As a summary of existing knowledge on the occurrence of *T. saginata* taeniosis and bovine cysticercosis in the area is presently lacking, and as part of a coordinated effort to document the global distribution of *T. saginata* [9–14], we undertook a systematic review of the occurrence of this parasite in West and Central Africa.

Methods

Search strategy

We conducted a systematic review aiming to gather current knowledge on the occurrence, prevalence and geographical distribution of human taeniosis and bovine cysticercosis in West and Central Africa, published between January 1st, 1990 and December 31st, 2017. A complete study protocol is available in Additional file 1: Text S1. In the context of this study, West and Central Africa was defined as the area covering the following 27 countries/territories: Ascension, Benin, Burkina Faso, Cameroon, Cape Verde, Central African Republic, Chad, Republic of the Congo, Democratic Republic of the Congo (DR Congo), Equatorial Guinea, Gabon, The Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Liberia, Mali, Mauritania, Niger, Nigeria, Saint Helena, Sao Tome and Principe, Senegal, Sierra Leone, Togo and Tristan da Cunha. Although Angola is classified as being part of Central Africa, it is also classified as being part of southern Africa, and data from this country were included in an equivalent systematic review of southern and eastern Africa [14].

The international scientific databases Web of Science (<http://ipsience.thomsonreuters.com/product/web-of-science/>) and PubMed (<http://www.ncbi.nlm.nih.gov/pubmed>) were searched using the following combination of key words: (cysticerc* OR cisticerc* OR “C. bovis” OR taenia* OR tenia* OR saginata OR taeniosis OR teniosis OR taeniasis OR ténia OR taeniid OR cysticercque) AND (Ascension OR Benin OR “Burkina Faso” OR Cameroon OR “Cape Verde” OR “Central African Republic” OR Chad OR Congo-Brazzaville OR DRC OR Congo OR “Cote d’Ivoire” OR “Equatorial Guinea” OR Gabon OR Gambia OR Ghana OR Guinea OR Guinée OR Guinea-Bissau OR Liberia OR Mali OR Mauritania OR Niger OR Nigeria OR “Saint Helena” OR Sao Tome OR Principe OR Senegal OR “Sierra Leone” OR Togo OR “Tristan da Cunha”). Moreover, databases for MSc/PhD theses and grey literature (Additional file 2: Text S2) were searched using the same search phrase. Furthermore, the OIE databases “Help with World Animal Disease Status” (“Handistatus”, 1996–2004) [15] and “World Animal Health Information System” (“WAHIS”, 2005) [16] were

consulted to extract data on bovine cysticercosis for the study area. Finally, reference lists of relevant reviews were screened for additional records.

Selection criteria

The PRISMA guidelines were followed for reporting the review (Additional file 3: Table S1). Briefly, duplicate records were removed after compiling results from the different searches, followed by screening of titles and abstracts for relevance. Then, full text articles were evaluated using the following exclusion criteria: (i) studies concerning a parasite different from *T. saginata*; (ii) studies conducted outside the study area; (iii) studies published outside the study period; (iv) studies reporting results outside the scope of our review question (e.g. review, experiment, intervention); and (v) duplicated data. No language restrictions were implemented.

Data extraction and compilation

Data from included records were extracted. Where records reported both the numerator and denominator of the study sample, respectively, prevalence and 95% Wilson score confidence intervals (CI) were calculated. All calculations were conducted in R, version 3.5.2.

Results

Search results

In total, 1670 records were retrieved, including 1655 through database searching, while 17 additional records were identified, including 15 retrieved through reference list screening, and the OIE databases Handistatus [15] and WAHIS [16]. After the removal of duplicates, out of 1237 remaining records, 1235 underwent title and abstract screening (i.e. the abstract was unavailable for 2 records). Subsequently, full texts of 87 articles were assessed for eligibility, of which 51 articles were retained for the qualitative synthesis (45 journal articles, 3 conference abstracts, 2 databases, 1 letter to the editor), of which 45 were included in the quantitative synthesis (Additional file 4: Figure S1).

Human taeniosis

A total of 45 records described human taeniosis cases, of which 39 were included in the quantitative synthesis (Table 1). Out of 45 records, 35 describe results from Nigeria, 3 from Cameroon, 3 from Senegal, 1 from Burkina Faso, 1 from DR Congo, 1 from Guinea and 1 from Ivory Coast (Fig. 1). No data were available for the other countries in the study area.

All studies included in the quantitative synthesis used plain stool microscopy as a diagnostic tool, and prevalence estimates for taeniosis ranged between 0–11% (0% in suburban schoolchildren [17] and 11% in community

residents [18] both in Nigeria). Three other studies performed in Nigeria; however, they reported much higher prevalence estimates: 23% in primary schoolchildren [19], 33% in rural schoolchildren [17] and even 50% in pre-school-age children (aged between 0 and 71 months) [20]. In four other studies, excluded from the quantitative synthesis, the presence of *T. saginata* taeniosis was described for Nigeria, Ivory Coast and Senegal, without prevalence estimates [21–24]. A further two case reports were also excluded from the qualitative synthesis: the first case described the presence of Meckel's diverticulitis due to *T. saginata* taeniosis in a 6-year-old girl in Nigeria [25], and the other a case of *T. saginata* taeniosis in a 33-year-old male presumably infected in Ivory Coast but diagnosed in Spain [26]. Only five studies reported the specific presence of *T. saginata* taeniosis; none of the studies mentioned morphological identification or molecular confirmation of the causative species, although one study conducted in Cameroon mentioned that species identification was done based on the expelled worm.

Bovine cysticercosis

Only 4 journal articles in addition to the 2 OIE databases described the presence of bovine cysticercosis in the study area, 3 of which were included in the quantitative synthesis (Table 2). The journal articles (1 described data for DR Congo, 3 for Nigeria) reported prevalence estimates based on abattoir surveys (i.e. meat inspection) between 0–29.0% (0% [27], 29% [28], both in Nigeria). One article, which was excluded from the quantitative synthesis, described the presence of bovine cysticercosis in Nigeria, without prevalence estimates [23].

In contrast with the journal articles, the OIE databases reported the (past) presence of bovine cysticercosis in a larger part of the study area (Table 3). Overall, bovine cysticercosis was reported throughout the study area, except for Guinea-Bissau, Sao Tomé and Príncipe, and Togo, where it was declared to be absent [15, 16] (Fig. 2). No data were available for Ascension, Cape Verde, the Republic of Congo, Equatorial Guinea, Gabon, Gambia, Liberia, Mauritania, Saint Helena or Sierra Leone.

Discussion

Our aim was to gather current knowledge on human taeniosis and bovine cysticercosis in West and Central Africa. Overall, human taeniosis was reported in seven out of 27 countries/territories, while bovine cysticercosis was reported in 14 out of 27 countries/territories. This systematic review has revealed that human *T. saginata* taeniosis and bovine cysticercosis are seriously understudied in this region. While the study area consists of 27 countries and territories, the presence of human taeniosis and bovine cysticercosis were only described for 7


Table 1 Reported occurrence of taeniosis in West and Central Africa

Country	Study period	n	n+	%	95% CI	Species	Groups studied	Reference
Burkina Faso	na	1587	na	2.1	na	N	Community volunteers	[39]
Cameroon	08/1999–04/2000	3109	1	0.03	0.006–0.18	Y ^a	Community volunteers	[35]
Cameroon	03/2012–07/2012	396	na	0.25	na	N	Patients consulting hospital	[40]
Cameroon	na	163	2	1.2	0.3–4.4	N	Pre-school-age children	[41]
DR Congo	03/04/2014–07/06/2018	602	40	6.6	4.9–8.9	Y ^b	School children (6–20 years)	[42]
Guinea	04/1995–06/1995	800	na	3.8	na	N	Children (10–14 years)	[43]
Nigeria	01/02/1997–31/01/1998	816	8	1.0	0.50–1.9	N	Pregnant women	[44]
Nigeria	07/02/1998–31/12/1998	129	3	2.3	0.8–6.6	Y ^b	Patients with complaints of upper abdominal pain, tenderness and indigestion	[45]
Nigeria	03/2000–09/2000	500	16	3.2	2.0–5.1	N	Schoolchildren	[46]
Nigeria	1/11/2003–30/1/2004	60	2	3.3	0.9–11.4	Y ^b	Children (2–6 years) attending clinic	[47]
Nigeria	11/2004–02/2005	232	23	9.9	6.7–14.4	N	Primary schoolchildren	[48]
Nigeria	01/2005–05/2005	309	5	1.6	[0.7; 3.7]	N	Schoolchildren	[49]
Nigeria	06/2005–11/2006	1059	102	9.6	8.0–11.6	N	Schoolchildren	[50]
Nigeria	10/2005–03/2006	119	1	0.8	0.2–4.6	N	Outpatients health institutions	[51]
Nigeria	1/2006–09/2006	73	24	32.9	23.2–44.3	N	Rural schoolchildren	[17]
Nigeria	1/2006–09/2006	171	19	11.1	7.2–16.7	N	Suburban schoolchildren	[17]
Nigeria	01/2006–09/2006	283	na	23.0	na	N	Primary schoolchildren	[19]
Nigeria	06/2006–11/2006	818	4	0.5	0.2–1.3	N	Hospital patients	[52]
Nigeria	01/2007–03/2007	250	1	0.4	0.07–2.2	N	Schoolchildren	[53]
Nigeria	07/2007–08/2007	100	13	13.0	7.8–21.0	N	Sampling in hostels (faeces sampling in toilets, not from individual participants)	[54]
Nigeria	08/2007–08/2009	500	0	0	0–0.8	N	HIV-negative enrolled via HIV outreach programme (in houses and offices)	[18]
Nigeria	08/2007–08/2009	2000	4	0.2	0.08–0.5	N	HIV-positive patients attending clinic	[18]
Nigeria	01/2008–08/2008	50	4	8	3.2–18.8	N	Villagers	[32]
Nigeria	05/2009–07/2009	122	2	1.6	0.5–5.8	N	Abattoir workers	[55]
Nigeria	05/2009–07/2009	98	na	0	na	N	Control populations	[55]
Nigeria	08/2010–12/2010	600	19	3.2	2.0–4.9	N	Schoolchildren	[56]
Nigeria	01/2011–12/2011	3826	na	0.89	na	N	Primary schoolchildren	[57]
Nigeria	06/2011–11/2011	220	na	5	na	N	Schoolchildren aged 1–15 years	[58]
Nigeria	06/2012–12/2012	116	10	8.6	4.7–15.1	Y ^b	Food vendors	[59]
Nigeria	09/2012–01/2013	717	5	0.7	0.3–1.6	N	Samples from polio surveillance programme	[60]
Nigeria	02/2014–06/2014	167	84	50.3	42.8–57.8	N	Pre-school-age children	[20]
Nigeria	07/2014–11/2014	112	3	2.7	0.9–7.6	N	Community volunteers	[61]
Nigeria	na	750	16	2.1	1.3–3.4	N	Women (child up to senior)	[62]
Nigeria	na	471	1	0.2	0.04–1.2	N	Primary schoolchildren	[63]
Nigeria	na	96	1	1	0.2–5.7	N	HIV patients	[64]
Nigeria	na	168	8	4.8	2.4–9.1	N	Food vendors	[65]
Nigeria	na	296	11	3.7	2.1–6.5	N	Villagers	[66]
Nigeria	na	162	2	1.2	0.3–4.4	N	Primary schoolchildren	[67]
Nigeria	na	400	12	3.0	1.7–5.2	N	Primary schoolchildren	[68]
Nigeria	na	416	45	10.8	8.2–14.2	N	Nursery and primary school children	[69]
Senegal	04/1997	400	na	3.5	na	N	Schoolchildren	[70]
Senegal	2004	na	na	4.6	na	N	Routine analyses at parasitology laboratory	[71]
Senegal	2005	na	na	5.6	na	N	Routine analyses at parasitology laboratory	[71]
Senegal	2006	na	na	4.0	na	N	Routine analyses at parasitology laboratory	[71]
Senegal	2007	na	na	7.0	na	N	Routine analyses at parasitology laboratory	[71]
Senegal	2008	na	na	5.5	na	N	Routine analyses at parasitology laboratory	[71]
Senegal	2009	na	na	4.1	na	N	Routine analyses at parasitology laboratory	[71]

^a Species identification based on expelled worm

^b Reported as *Taenia saginata*, yet unclear from methodology how species identification was done

Abbreviations: n, number of individuals tested; n+, number of positive individuals; CI, confidence interval; na, not available; Y, yes; N, no



and 17 countries, respectively. For the remaining regions no data were reported. These findings are in contrast to eastern and southern Africa which have similar lifestyles yet a remarkably higher coverage [12]. This difference might be partially explained by the presence of a large French speaking population in West and Central Africa, and therefore potentially published research in journals which were not detected completely by our search strategy, although we had included French search terms and some articles in French were included in the present study. Another explanation might be a lower awareness and interest in the topic, as the cattle population in this area is somewhat lower as compared to eastern and southern Africa, i.e. 120 million heads [7] as opposed to 184 million heads [7].

Overall, the reported prevalence estimates of taeniosis were in line with those reported in eastern and southern Africa [12] and in the Americas [13], but higher than

those reported in western and eastern Europe [9, 10]. In contrast to certain areas in eastern and southern Africa where consumption of raw beef is a culinary habit [12], traditional dishes in West and Central Africa include mainly stews with typically extended cooking times [29, 30], thereby decreasing the risk of exposure to viable *T. saginata* cysts. Three articles, however, reported very high taeniosis prevalence estimates, with 23% in primary schoolchildren [19], 33% in rural schoolchildren [17] and even 50% in pre-school-age children [20]. According to Adeniran et al. [20], Nigerian pre-school-age children are often fed undercooked meals for adults, including beef, in order to facilitate the transition from breastmilk to solid food. Should this high prevalence estimate be confirmed, such practices should be investigated and be the subject of close attention and education on the potential dangers associated with consumption of inadequately cooked food. In all taeniosis cases, species identification

Table 2 Reported occurrence of bovine cysticercosis in West and Central Africa: reports based on meat inspection


Country	Study period	Data source	<i>n</i>	<i>n</i> +	%	95% CI	Reference
Congo	06/1986–06/1987	Veterinary inspection records	3914	na	9.6	na	[72]
Congo	07/1986–10/1988	Veterinary inspection records	333	na	10.5	na	[72]
Congo	07/1986–10/1988	Veterinary inspection records	284	na	12.7	na	[72]
Congo	na	Veterinary inspection records	73	na	1.4	na	[72]
Congo	na	Veterinary inspection records	47	na	4.3	na	[72]
Congo	na	Veterinary inspection records	35	na	14.3	na	[72]
Nigeria	1985–1986	Veterinary inspection records	1221	0	0	0–0.3	[27]
Nigeria	1985–1986	Retail market inspection by investigators	358	27	7.5	5.2–10.8	[27]
Nigeria	11/1999–04/2002	Carcass inspection by investigators	5560	1,610	29.0	27.8–30.2	[28] ^a
Nigeria	11/1999–04/2002	Carcass inspection by investigators	20,240	5,140	25.4	24.8–26.0	[28] ^b
Nigeria	2005–2007	Veterinary inspection records	641,224	805	0.13	0.12–0.13	[73]

^a Local breeds in rural areas^b Exotic breeds in urban areasAbbreviations: *n*, number of individuals tested; *n*+, number of positive individuals; CI, confidence interval; na, not available**Table 3** OIE data on occurrence of bovine cysticercosis in West and Central Africa (1996–2005) [15, 16]

Country/territory	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Ascension										
Benin								+	79	
Burkina Faso	+	+	+							+
Cameroon		+	+	+	+	+	+	21	23	+
Cape Verde										
Central African Republic	+					+	+		+	+
Chad							+			
DR Congo							+			+
Republic of the Congo										
Equatorial Guinea										
Gabon										
Gambia										
Ghana		+								-
Guinea	+									
Guinea-Bissau							-	-	-	
Ivory Coast	+	+	+	+	+	+	+	+	+	
Liberia										
Mali	+									-
Mauritania										
Niger	+	+				+				
Nigeria				+		+	+	153	34	-
Saint Helena										
Sao Tome and Principe					-	-	-	-	-	
Senegal	92	+	199	125	+	+				
Sierra Leone										
Togo										-
Tristan da Cunha		71			1	+		1		

Blank cells indicate the data were unavailable

Abbreviations: CI, confidence interval; +, occurrence of the disease; -, absence of the disease



is pivotal to avoid the potential ingestion of *T. solium* eggs by the tapeworm carrier, and relatives and acquaintances, representing a risk of neurocysticercosis. *Taenia solium* is endemic in many of the included countries or territories [31–36]. Most studies, however, reported non-specified taeniosis, and for those specifically mentioning *T. saginata* taeniosis only one described the applied method. None of the other studies reported the use of specific morphological methods or molecular tools allowing for the identification of the causative *Taenia* sp. [9]. Hence, we cannot rule out that a certain proportion of taeniosis cases summarized in this review are due to *T. solium* instead of *T. saginata*.

A wide fluctuation in prevalence estimates, ranging between 10–30% for most studies [37] was observed for bovine cysticercosis, which is comparable to the estimates for eastern and southern Africa [12]. In many

countries, meat inspection is not done systematically, especially in the case of backyard slaughtering in rural areas where meat inspection may not be available. Moreover, a correct estimation of the prevalence through meat inspection is hampered by its low sensitivity, which is estimated to lie below 16% [38]. This was confirmed by one study from Nigeria describing the prevalence of bovine cysticercosis in carcasses having passed the regular meat inspection at the abattoir. While the regular veterinary inspection declared the carcasses to be free from cysticercosis, investigators found a prevalence of 7.5% in carcasses originating from the same abattoir, sold at retail markets [27]. Overall, prevalence estimates for taeniosis of up to 50%, and for bovine cysticercosis of up to 30%, indicate the continued transmission of this parasite between cattle and humans. Despite the limited pathology caused by *T. saginata*, bovine cysticercosis has the

potential to cause a high economic cost due to condemnation of infected carcasses. To interrupt transmission, stringent meat inspection procedures should be applied and improvements implemented in the sanitation and management of human sewage. The public should also be educated regarding general food safety measures such as thorough cooking of meat products, which also reduces the risk of infection with other microbiological hazards associated with meat products, such as pathogenic *Escherichia coli*, *Salmonella* spp., *Mycobacterium bovis* and *Campylobacter* spp.

Conclusions

Based on the findings of our systematic review, both human taeniosis and bovine cysticercosis are understudied in West and Central Africa. Included articles reported high prevalence estimates for both conditions, pointing to a continued transmission of *T. saginata* in the region. A One Health approach is needed to protect the general public from acquiring tapeworm infection.

Additional files

- Additional file 1: Text S1.** Search protocol.
- Additional file 2: Text S2.** Databases used.
- Additional file 3: Table S1.** PRISMA checklist.
- Additional file 4: Figure S1.** PRISMA flow diagram.

Abbreviations

CI: confidence interval; DR Congo: Democratic Republic of the Congo; OIE: World Organisation for Animal Health/Office International des Epizooties; PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; WAHIS: World Animal Health Information System.

Acknowledgements

This work was performed within the framework of CYSTINET, the European network on taeniosis/cysticercosis, COST ACTION TD1302.

Authors' contributions

EH and VD conducted the systematic review of literature, extracted and analysed the data, and drafted the first version of the manuscript. All authors contributed to the design of the study, interpretation of the data and writing of the paper. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

All references found eligible in our literature review are included in the article.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium. ² International Livestock Research Institute (ILRI), P.O. Box 30709, Nairobi, Kenya. ³ Institute for Infection & Global Health, University of Liverpool, Neston, UK. ⁴ Department of Virology, Parasitology and Immunology, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium. ⁵ Centre of Excellence for Food and Vector-borne Zoonoses, Institute for Medical Research, University of Belgrade, Dr Subotića 4, Belgrade 11000, Serbia. ⁶ One Health Center for Zoonoses and Tropical Veterinary Medicine, Ross University School of Veterinary Medicine, P.O. Box 334, Basseterre, Saint Kitts and Nevis. ⁷ Department of Infectious Disease Epidemiology and Prevention, Statens Serum Institut, 2300 Copenhagen, Denmark. ⁸ Department of Veterinary Public Health and Food Safety, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium. ⁹ Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium. ¹⁰ Institute of Parasitology, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland. ¹¹ Veterinary Research Institute, Hellenic Agricultural Organisation Demeter, Thessaloniki 57001, Greece. ¹² Department of Food Safety and Infection Biology, Faculty of Veterinary Medicine, Norwegian University of Life Sciences, Oslo, Norway.

Received: 8 April 2019 Accepted: 24 June 2019

Published online: 27 June 2019

References

1. Craig P, Ito A. Intestinal cestodes. *Curr Opin Infect Dis*. 2007;20:524–32.
2. Tembo A, Craig PS. *Taenia saginata* taeniosis: copro-antigen time-course in a voluntary self-infection. *J Helminthol*. 2015;89:612–9.
3. Gonzalez A, Thomas L. *Taenia* spp. In: Rose JB, Jiménez-Cisneros B, editors. *Global Water Pathogen Project*. 2018. <http://www.waterpathogens.org>. Accessed 21 Jun 2019.
4. Symeonidou I, Arsenopoulos K, Tzilves D, Soba B, Gabriël S, Papadopoulos E. Human taeniosis/cysticercosis: a potentially emerging parasitic disease in Europe. *Ann Gastroenterol*. 2018;31:406–12.
5. Blagojevic B, Robertson LJ, Vieira-Pinto M, Vang Johansen M, Laranjo-González M, Gabriël S. Bovine cysticercosis in the European Union: impact and current regulations, and an approach towards risk-based control. *Food Control*. 2017;78:64–71.
6. Jansen F, Dorny P, Trevisan C, Dermauw V, Laranjo-González M, Allepuz A, et al. Economic impact of bovine cysticercosis and taeniosis caused by *Taenia saginata* in Belgium. *Parasit Vectors*. 2018;11:241.
7. Food and Agriculture Organization of the United Nations statistical databases. Rome: FAO; 2019. <http://faostat3.fao.org>. Accessed 21 Jun 2019.
8. Otte M, Chilonda P. Cattle and small ruminant production systems in sub-Saharan Africa—a systematic review. Rome: Food and Agriculture Organization of the United Nations; 2002.
9. Laranjo-González M, Devleeschauwer B, Trevisan C, Allepuz A, Sotiraki S, Abraham A, et al. Epidemiology of taeniosis/cysticercosis in Europe, a systematic review: western Europe. *Parasit Vectors*. 2017;10:349.
10. Trevisan C, Sotiraki S, Laranjo-González M, Dermauw V, Wang Z, Kärssin A, et al. Epidemiology of taeniosis/cysticercosis in Europe, a systematic review: eastern Europe. *Parasit Vectors*. 2018;11:569.
11. Bobić B, Thomas LF, Djurković OD, Devleeschauwer B, Dermauw V, Dorny P, et al. Epidemiology of *Taenia saginata* taeniosis/cysticercosis in the Russian Federation. *Parasit Vectors*. 2018;11:636.
12. Saratsis A, Sotiraki S, Braae UC, Devleeschauwer B, Dermauw V, Eichenberger RM, et al. Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in the Middle East and North Africa. *Parasit Vectors*. 2019;12:113.
13. Braae UC, Thomas LF, Robertson LJ, Dermauw V, Dorny P, Willingham AL, et al. Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in the Americas. *Parasit Vectors*. 2018;11:518.
14. Dermauw V, Dorny P, Braae UC, Devleeschauwer B, Robertson LJ, Saratsis A, et al. Epidemiology of *Taenia saginata* taeniosis/cysticercosis: a systematic review of the distribution in the southern and eastern Africa. *Parasit Vectors*. 2018;11:578.

15. OIE Handistatus II. Office International des Epizooties, Paris. 2018. <http://web.oie.int/hs2/report.asp>. Accessed 8 Apr 2019.
16. OIE World Animal Health Information Database (WAHIS). Office International des Epizooties, Paris. 2018. http://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home. Accessed 8 Apr 2019.
17. Auta T, Reuben WJ, Abdulhadi JB, Mohammed JA. A comparative study on the prevalence of intestinal helminthes among rural and sub-urban pupils in Gwagwada, Nigeria. *J Parasitol Vector Biol*. 2014;5:87–91.
18. Akinbo FO, Okaka CE, Omoregie R. Prevalence of intestinal parasitic infections among HIV patients in Benin City, Nigeria. *Libyan J Med*. 2010;5:5506.
19. Auta T, Kogi E, Kokori Audu O. Studies on the intestinal helminths infestation among primary school children in Gwagwada, Kaduna, North Western Nigeria. *J Biol Agric Healthc*. 2013;3:48–54.
20. Adeniran AA, Mogaji HO, Aladesida AA, Olayiwola IO, Oluwole AS, Abe EM, et al. Schistosomiasis, intestinal helminthiasis and nutritional status among preschool-aged children in sub-urban communities of Abeokuta, Southwest, Nigeria. *BMC Res Notes*. 2017;10:637.
21. Adoke K, Iliyasu Y, Adoke A, Ahmed S. Neglected tropical diseases: A histopathological review. *Virchows Arch*. 2017;471:530.
22. Ademowo GO, Rabiou O, Kosoko A, Dada-adegbola H, Arinola G, Falade C. Prevalence and interaction of malaria and helminth co-infections among symptomatic and asymptomatic children in Southwest Nigeria. *Int J Infect Dis*. 2014;21:297.
23. Ekong PS, Juryit R, Dika NM, Nguku P, Musenero M. Prevalence and risk factors for zoonotic helminth infection among humans and animals—Jos, Nigeria, 2005–2009. *Pan Afr Med J*. 2012;12:2005–9.
24. Secka A, Grimm F, Marcotty T, Geysen D, Niang AM, Ngale V, et al. Old focus of cysticercosis in a senegalese village revisited after half a century. *Acta Trop*. 2011;119:199–202.
25. Chirdan L, Yusufu L, Ameh E, Shehu S. Meckel's diverticulitis due to *Taenia saginata*: a case report. *East Afr Med J*. 2001;78:107–8.
26. Lopez-Caleya JF, Contreras SN, Martin-Rodrigo L. *Taenia saginata*: an imported case. *Rev Esp Enferm Dig*. 2015;107:440–1.
27. Ogunrinade A, Oyekole O. Evaluation of the efficiency of beef inspection procedures for tuberculosis, fascioliasis and cysticercosis in a Nigerian abattoir. *Prev Vet Med*. 1990;8:71–5.
28. Opara MN, Ukpong UM, Okoli IC, Anosike JC. Cysticercosis of slaughter cattle in southeastern Nigeria. *Ann NY Acad Sci*. 2006;1081:339–46.
29. Jacob J, Ashkenazi M. The world cookbook: the greatest recipes from around the globe. Santa Barbara: Greenwood; 2014.
30. Massaquoi R. Foods of Sierra Leone and other West African countries, a cookbook and food-related stories. Bloomington: Authorhouse; 2011.
31. Madinga J, Kanobana K, Lukanu P, Abatih E, Baloji S, Linsuke S, et al. Geospatial and age-related patterns of *Taenia solium* taeniasis in the rural health zone of Kimpese, Democratic Republic of Congo. *Acta Trop*. 2016;165:100–9.
32. Gweba M, Faleke OO, Junaidu A, Fabiya JP, Fajinmi AO. Some risk factors for *Taenia solium* cysticercosis in semi-intensively raised pigs in Zuru, Nigeria. *Vet Ital*. 2010;46:57–67.
33. Dermauw V, Carabin H, Ganaba R, Cissé A, Tarnagda Z, Gabriël S, et al. Factors associated with the 18-month cumulative incidence of sero-conversion of active infection with *Taenia solium* cysticercosis: a cohort study among residents of 60 villages in Burkina Faso. *Am J Trop Med Hyg*. 2018;99:1018–27.
34. Permin A, Yelifari L, Bloch P, Steenhard N, Hansen NP, Nansen P. Parasites in cross-bred pigs in the Upper East Region of Ghana. *Vet Parasitol*. 1999;87:63–71.
35. Vondou L, Zoli AP, Nguekam, Pouedet S, Assana E, Kamga Tokam AC, et al. La taeniose/cysticercose à *Taenia solium* dans la Menoua (Ouest-Cameroun). *Parasite*. 2002;9:271–4.
36. Braae UC, Saarnak CFL, Mukaratirwa S, Devleeschauwer B, Magnussen P, Johansen MV. *Taenia solium* taeniosis/cysticercosis and the co-distribution with schistosomiasis in Africa. *Parasit Vectors*. 2015;8:323.
37. Laranjo-González M, Devleeschauwer B, Gabriël S, Dorny P, Allepuz A. Epidemiology, impact and control of bovine cysticercosis in Europe: a systematic review. *Parasit Vectors*. 2016;9:81.
38. Eichenberger RM, Lewis F, Gabriël S, Dorny P, Torgerson PR, Deplazes P. Multi-test analysis and model-based estimation of the prevalence of *Taenia saginata* cysticercus infection in naturally infected dairy cows in the absence of a "gold standard" reference test. *Int J Parasitol*. 2013;43:853–9.
39. Ouedraogo A, Tiono A, Amidou D, Ouedraogo A, Ouedraogo E, Kabore Y, et al. Site characterization for a malaria vaccine trial in the Saponé health district in Burkina Faso: seasonal prevalence of main parasites infestation. *Am J Trop Med Hyg*. 2010;83:48.
40. Nkenfou CN, Nana CT, Payne VK. Intestinal parasitic infections in HIV infected and non-infected patients in a low HIV prevalence region, West-Cameroon. *PLoS ONE*. 2013;8:e57914.
41. Ajeagah G, Wouafo M, Ezenguele G, Nzukam J. Presence of gastrointestinal parasites in a tropical urban region (Yaoundé, Cameroon). *Comp Parasitol*. 2013;80:279–83.
42. Kyambikwa Bisangamo C, Jabari Mutwa P, Mulongo Mbarambara P. Profil des parasitoses intestinales chez les enfants d'âge scolaire de Kiliba (est de la RD Congo). *Med Sante Trop*. 2017;27:209–13.
43. Gyorkos TW, Camara B, Kokoskin E, Carabin H, Prouty R. Enquête de prévalence parasitaire chez les enfants d'âge scolaire en Guinée (1995). *Sante*. 1996;6:377–81.
44. Egwunyenga AO, Ajayi JA, Nmorsi OPG, Duhlinska-Popova DD. *Plasmodium*/intestinal helminth co-infections among pregnant Nigerian women. *Mem Inst Oswaldo Cruz*. 2001;96:1055–9.
45. Anyaeze CM. Reducing burden of hookworm disease in the management of upper abdominal pain in the tropics. *Trop Doct*. 2003;33:174–5.
46. Kamalu N, Uwakwe F, Opara J. Prevalence of intestinal parasite among high school students in Nigeria. *Acad J Interdiscip Stud*. 2013;2:9–16.
47. Okeniyi JAO, Ogunlesi TA, Oyelami OA, Adeyemi LA. Effectiveness of dried carica papaya seeds against human intestinal parasitosis: a pilot study. *J Med Food*. 2007;10:194–6.
48. Ekpo UF, Odoemene SN, Mafiana CF, Sam-Wobo SO. Helminthiasis and hygiene conditions of schools in Ikenne, Ogun State, Nigeria. *PLoS Negl Trop Dis*. 2008;2:e146.
49. Ugbomoiko US, Ofioezie IE. Multiple infection diagnosis of intestinal helminthiasis in the assessment of health and environmental effect of development projects in Nigeria. *J Helminthol*. 2007;81:227–31.
50. Agbolade OM, Agu NC, Adesanya OO, Odejayi AO, Adigun AA, Adeslanu EB, et al. Intestinal helminthiasis and schistosomiasis among school children in an urban center and some rural communities in southwest Nigeria. *Korean J Parasitol*. 2007;45:233.
51. Abaver DT, Nwobegahay JM, Goon DT, Iweriebor BC, Anye DN. Prevalence of intestinal parasitic infections among HIV/AIDS patients from two health institutions in Abuja, Nigeria. *Afr Health Sci*. 2011;11:524–7.
52. Abelau M, Goselle O, Udeh E, D-Popova D, Popov T. A comparative study of the prevalence of helminth parasites in HIV seropositive and HIV seronegative individuals in plateau state. *Niger Ann Nat Sci*. 2011;11:22–32.
53. Damen JG, Lar P, Mershak P, Mbaawuga EM, Nyary BW. A comparative study on the prevalence of intestinal helminthes in dewormed and non-dewormed students in a rural area of North-Central Nigeria. *Lab Med*. 2010;41:585–9.
54. Ejima L, Ajogun R. The prevalence and health implications of the ova of human intestinal helminth parasites isolated from faeces collected near students' hostels, federal polytechnic, Idah, Kogi State, Nigeria. *Int J Trop Med*. 2011;6:15–8.
55. Banjo T, Amoo A, Busari A, Kama A, Lawal I, Ogundahunsi O, et al. Intestinal parasites among abattoir workers in Abeokuta. *Am J Res Commun*. 2013;1:84–96.
56. Shehu M, Kabiru A, Abubakar U, Muhammad K. prevalence of intestinal helminth infections among school children in relation to occupation of parents and toilets facilities in Maru LGA Zamfara State. *J Biol Agric Healthc*. 2013;3:87–91.
57. Abah AE, Arene FOI. Status of intestinal parasitic infections among primary school children in Rivers State, Nigeria. *J Parasitol Res*. 2015;2015:937096.
58. Nwalorzie C, Onyenakazi S, Ogwu S, Okafor A. Predictors of intestinal helminthic infections among school children in Gwagwalada, Abuja, Nigeria. *Niger J Med J Natl Assoc Resid Dr Niger*. 2015;24:233–41.
59. Omalu ICJ, Paul S, Adeniran LA, Hassan SC, Pam VA, Eke SS, et al. Assessment of the level of gastrointestinal parasites infection among food vendors in Minna, North central Nigeria. *Annu Rev Res Biol*. 2013;3:705–13.
60. Adekolujou DR, Olayinka SO, Adeniji JA, Oyejemi OT, Odaibo AB. Poliovirus and other enteroviruses in children infected with intestinal parasites in Nigeria. *J Infect Dev Ctries*. 2015;9:1166–71.
61. Taiwo OT, Sam-Wobo SO, Idowu OA, Talabi AO, Taiwo AM. Comparative assessment of intestinal helminths prevalence in Water, Sanitation and

- Hygiene (WASH) intervention and non-intervention communities in Abeokuta, Nigeria. *Asian Pac J Trop Biomed*. 2017;7:524–32.
62. Amuta E, Houmsou R, Mker S. Knowledge and risk factors of intestinal parasitic infections among women in Makurdi, Benue State. *Asian Pac J Trop Med*. 2010;3:993–6.
63. Dangana A, Abayomi R, Way G, Akobi O. Survey of *Ascaris lumbricoides* among pupils of primary school in Jos south local government area of Plateau State, Nigeria. *Afr J Microbiol Res*. 2011;5:2524–7.
64. Ojurongbe O, Raji OA, Akindele AA, Kareem MI, Adefioye OA, Adeyeba AO. *Cryptosporidium* and other enteric parasitic infections in HIV-seropositive individuals with and without diarrhoea in Osogbo, Nigeria. *Br J Biomed Sci*. 2011;68:75–8.
65. Ifeadike C, Ironkwe O, Adogu POU, Nnebue C, Emelumadu O, Nwabueze S, et al. Prevalence and pattern of bacteria and intestinal parasites among food handlers in the Federal Capital Territory of Nigeria. *Niger Med J*. 2012;53:166–71.
66. Enimien OJ, Fana SA, Emmanuel WB. Intestinal helminthic infection in Numan (Northeast Nigeria). *Int J Progress Sci Technol*. 2015;1:1–4.
67. Ojurongbe O, Oyesiji K, Ojo J, Odewale G, Adefioye O, Olowe A, et al. Soil transmitted helminth infections among primary school children in Ile-Ife Southwest, Nigeria: a cross-sectional study. *Int Res J Med Med Sci*. 2014;2:6–10.
68. Thomas H, Jatau E, Inabo H, Garba D. Prevalence of intestinal helminths among primary school children in Chikun and Kaduna South Local Government areas of Kaduna state, Nigeria. *J Med Med Res*. 2014;2:6–11.
69. Umeh C, Mbanugo J, Ezeugoigwe N. Prevalence of intestinal helminth parasite in stools of nursery and primary schools pupils in Uga, Anambra State, Nigeria. *Sky J Microbiol Res*. 2015;3:6–10.
70. Diouf S, Diallo A, Camara B, Diagne I, Ndiaye O, Tall A, et al. Etat vaccinal et pathologies de l'enfant en zone rurale Senegalaise (Khombole). *Arch Pediatr*. 1999;6:903.
71. Ndiaye D, Ndiaye M, Gueye PAL, Badiane A, Fall ID, Ndiaye YD, et al. Prévalence des helminthoses digestives diagnostiquées à l'hôpital Le Dantec de Dakar, Sénégal. *Med Sante Trop*. 2013;23:35–8.
72. Chartier C, Bushu M, Kamwenga D. Les dominantes du parasitisme helminthique chez les bovins en Ituri (Haut-Zaïre). III. Répartition géographique et prévalence des principaux helminthes. *Rev Elev Med Vet Pays Trop*. 1991;44:61–8.
73. Cadmus SIB, Adesokan HK. Causes and implications of bovine organs/offal condemnations in some abattoirs in western Nigeria. *Trop Anim Health Prod*. 2009;41:1455–63.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

